



35% **PROGRESS: HeartCode ACLS 2025**

-4h 11m left

Ahmed Othman



## Coach



Learn more here:

☐ EMS Transport and Aler...



What is an advantage of EMS alerting the receiving facility of the impending arrival of a patient with suspected acute ischemic stroke?

You got it!



Your Answer

| The hospital can perform more efficient evaluation and management.



I Know It

CHALLENGE US

NEXT

## Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER




Coach



Push ► for reading content aloud.

Autoplay ☒ On



## ACUTE CORONARY SYNDROMES INTRODUCTION



### Goals for ACS Patients

The primary goals for ACS patients are

- Prevention of major adverse cardiovascular events such as death, nonfatal myocardial infarction (MI), and the need for urgent post-infarction revascularization.
- Identification of patients with STEMI and triage for early reperfusion therapy.
- Relief of ischemic chest discomfort.
- Treatment of acute, life-threatening complications of ACS, such as ventricular fibrillation (VF) or pulseless ventricular tachycardia (pVT), unstable bradycardias, ventricular wall rupture, papillary muscle rupture, decompensated shock, and other unstable tachycardias.

1

2

3

4

NEXT

CHALLENGE US





18%

PROGRESS: Hear

To exit full screen, move mouse to top of screen or press

F11

4h 7m left



Ahmed Otman

A




Coach



Push ► for reading content  
aloud.

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the  
right to focus on the questions



## ACUTE CORONARY SYNDROMES INTRODUCTION



### OHCA Response

Half of ACS deaths occur before the patient reaches the hospital, with VF or pVT as the precipitating rhythm in the majority of cases.

VF is most likely to develop during the first 4 hours after onset of symptoms, so communities should develop programs to respond quickly to ACS.

Such programs should focus on

- Recognizing symptoms of ACS.
- Activating the emergency medical services (EMS) system, with EMS providing prehospital notification in advance.
- Providing early CPR if cardiac arrest occurs.
- Providing early defibrillation with AEDs available through public-access defibrillation programs and first responders.
- Providing a coordinated system of care among the EMS system, the emergency department (ED), and cardiac specialists.

PREVIOUS

1

2

3

4

NEXT



## ACUTE CORONARY SYNDROMES INTRODUCTION

### STEMI Chain of Survival

The STEMI Chain of Survival is similar to the Chain of Survival for sudden cardiac arrest. Its links indicate the actions that patients, family members, and healthcare providers can rapidly take to maximize STEMI recovery:

- Recognition and reaction to STEMI warning signs.
- EMS dispatch and rapid EMS system transport and prearrival notification to the receiving hospital.
- Assessment and diagnosis in the ED (or cath lab).
- Treatment.



PREVIOUS

1

2

3

4

NEXT

CHALLENGE US



Coach



Push ► for reading content aloud.

Autoplay ☒ On



### Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



Coach



Push ► for reading content aloud.

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions




ADVANCED BEGINNER

# ACUTE CORONARY SYNDROMES INTRODUCTION

## ACS Algorithm

The Acute Coronary Syndromes (ACS) Algorithm will help guide your clinical strategy when patients have signs and symptoms of ACS, including possible acute myocardial infarction.

Acute Coronary Syndromes Algorithm



PREVIOUS 1 2 3 4

I KNEW GOT IT NOW THINK I GOT IT I DON'T GET IT

CHALLENGE US ?



## Coach



Push ► for reading content aloud.

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## SIGNS AND SYMPTOMS OF ACS

The most common symptom of myocardial ischemia and infarction is retrosternal chest discomfort. The patient may perceive this discomfort more as pressure or tightness than as actual pain.

Chest discomfort is the major symptom in most patients (both men and women) with ACS, but patients frequently deny or misinterpret this and other symptoms. The elderly, women, diabetic patients, and hypertensive patients are most likely to delay, in part because they are more likely to have atypical symptoms or presentations.

Keep in mind that older adults and women may present without chest pain.

## Symptoms that suggest ACS may also include

- Uncomfortable pressure, fullness, squeezing, or pain in the center of the chest lasting several minutes (usually more than a few minutes)
- Chest discomfort spreading to the shoulders, neck, one or both arms, or jaw
- Chest discomfort spreading into the back or between the shoulder blades
- Light-headedness, dizziness, fainting, sweating, nausea, or vomiting

I KNEW

GOT IT NOW

THINK I GOT IT

I DON'T GET IT

CHALLENGE US







18%

PROGRESS: HeartCode ACLS 2025

~4h 5m left



Ahmed Othman



Coach



Push ► for reading content  
aloud.

Autoplay ☒ On

Self-Assessment ?

Adjust your competence estimate to the  
right to focus on the questions



ADVANCED BEGINNER

## SIGNS AND SYMPTOMS OF ACS

Keep in mind that older adults and women may present without chest pain.

### Symptoms that suggest ACS may also include

- Uncomfortable pressure, fullness, squeezing, or pain in the center of the chest lasting several minutes (usually more than a few minutes)
- Chest discomfort spreading to the shoulders, neck, one or both arms, or jaw
- Chest discomfort spreading into the back or between the shoulder blades
- Light-headedness, dizziness, fainting, syncope, sweating, nausea, or vomiting
- Unexplained sudden shortness of breath, which may occur with or without chest discomfort
- Less commonly, the discomfort occurs in the epigastrium and is described as indigestion

These symptoms may also suggest other life-threatening conditions, including aortic dissection, acute pulmonary embolism, acute pericardial effusion with tamponade, and tension pneumothorax.

I KNEW

GOT IT NOW

THINK I GOT IT

I DON'T GET IT

CHALLENGE US





19%

PROGRESS: HeartCode ACLS 2025

4h 4m left

Ahmed Othman



## Coach



Learn more here:

Signs and Symptoms of ...



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

What is the most common symptom of myocardial ischemia and infarction?

You got it!



Your Answer

Retrosternal chest pain



I Know It

CHALLENGE US

NEXT





Coach



Learn more here:

Signs and Symptoms of ...



Which demographic group experiencing acute coronary syndromes is more likely to present without chest pain?

You got it!



Your Answer

Females

Older adults may also present without chest pain.



I Know It

CHALLENGE US

NEXT

Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



20%

PROGRESS: HeartCode ACLS 2025

~4h 5m left



Ahmed Othman


A



Coach



Push ► for reading content aloud.

Autoplay ☒ On

## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## EMS ASSESSMENT, CARE, AND HOSPITAL PREPARATION

Step 2 in the algorithm outlines EMS assessment, care, and hospital preparation. EMS responders may perform the following assessments and actions as they stabilize, triage, and transport the patient to an appropriate facility.

- **Assess ABC** (airway, breathing, circulation). Be prepared to provide CPR and defibrillation.
- **Administer aspirin** and **consider oxygen, nitroglycerin, and morphine** if needed.
- **Obtain a 12-lead ECG**. If there is ST elevation, notify the receiving hospital with a transmission or interpretation; note the time of onset and first medical contact.
- **Provide prehospital notification**; on arrival, transport to ED/cath lab per protocol.
- The notified hospital should mobilize resources to respond to STEMI.
- If considering prehospital fibrinolysis, **use a fibrinolytic checklist**.
- If out-of-hospital providers cannot complete these initial steps before the patient arrives at the hospital, the ED provider should do so.



Obtaining a 12-Lead ECG

I KNEW

GOT IT NOW

THINK I GOT IT

I DON'T GET IT

CHALLENGE US






## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## EMS ASSESSMENT, CARE, AND HOSPITAL PREPARATION

- Administer aspirin and consider oxygen, nitroglycerin, and morphine if needed.
- Obtain a 12-lead ECG. If there is ST elevation, notify the receiving hospital with a transmission or interpretation, note the time of onset and first medical contact.
- Provide prehospital notification: on arrival, transport to ED/cath lab per protocol.
- The notified hospital should mobilize resources to respond to STEMI.
- If considering prehospital fibrinolysis, use a fibrinolytic checklist.
- If out-of-hospital providers cannot complete these initial steps before the patient arrives at the hospital, the ED provider should do so.



### Obtaining a 12-Lead ECG

The American Heart Association (AHA) recommends out-of-hospital 12-lead ECG diagnostic programs in all EMS systems, and all EMS systems should take the actions outlined in the table.

EMS actions per AHA recommendations: [\[link\]](#)



CHALLENGE US ?

Push ► for reading content

contact

### EMS Action

### Recommendation

Obtain a 12-lead ECG if available

The AHA recommends routine use of 12-lead out-of-hospital ECGs for patients with signs and symptoms of possible ACS.

Provide prearrival notification to the hospital

Preambival notification of the ED shortens the time to treatment (10 to 60 minutes has been achieved in clinical studies) and speeds reperfusion therapy with fibrinolytics or PCI or both, which may reduce mortality and minimize myocardial injury.

Complete a fibrinolytic checklist if appropriate

If STEMI is identified on the 12-lead ECG, complete a fibrinolytic checklist if appropriate.

CLOSE

## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions.

EMS actions per AHA recommendations



I KNEW

GOT IT NOW

THINK I GOT IT

I DON'T GET IT



20% PROGRESS HeartCode ACLS 2025

4h 6m left

Ahmed Othman



Coach



Push ▶ for reading content aloud

Autoplay ☒ On



I Know It

CHALLENGE US



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



20% **PROGRESS** HeartCode ACLS 2025

4h 6m 10s

Ahmed Othman



## Coach



Learn more here:

☐ EMS Assessment, Care...



## Self-Assessment

Adjust your competence estimate to the right to focus on the questions.



ADVANCED BEGINNER

Obtaining a  is the most important assessment tool for a patient displaying signs and symptoms of acute coronary syndrome.

You got it!



Know it

CHALLENGE US

**NEXT**





## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

# CONCURRENT ED OR CATH LAB ASSESSMENT

The ED high-performance team should quickly evaluate the patient with potential ACS on the patient's arrival. Within the first 10 minutes, obtain a 12-lead ECG (if not already performed before arrival) and assess the patient.

### Initial Actions

- Activate STEMI team upon EMS notification.
- Assess ABCs: give oxygen if needed
- Check vital signs and evaluate oxygen saturation.
- Establish intravenous (IV) access
- Perform a brief, targeted history and a physical exam.
- Review and complete the fibrinolytic checklist: check contraindications.
- Obtain initial cardiac marker levels, complete blood counts, and coagulation studies
- Obtain a portable chest x-ray (in less than 30 minutes): do not delay transport to the cath lab

1

2

3

NEXT

CHALLENGE US





## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## CONCURRENT ED OR CATH LAB ASSESSMENT

### Initial Actions

- Activate STEMI team upon EMS notification.
- Assess ABCs: give oxygen if needed.
- Check vital signs and evaluate oxygen saturation.
- Establish intravenous (IV) access.
- Perform a brief, targeted history and a physical exam.
- Review and complete the fibrinolytic checklist: check contraindications.
- Obtain initial cardiac marker levels, complete blood counts, and coagulation studies.
- Obtain a portable chest x-ray (in less than 30 minutes): **do not delay transport to the cath lab**.

The results of cardiac markers, chest x-ray, and laboratory studies should not delay reperfusion therapy unless clinically necessary, for example, in suspected aortic dissection or coagulopathy.

1

2

3

NEXT

CHALLENGE US ?




Coach



Push ▶ for reading content aloud

Autoplay ☒ On



Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## CONCURRENT ED OR CATH LAB ASSESSMENT

Perform a targeted evaluation focusing on chest discomfort, signs and symptoms of heart failure, cardiac history, risk factors for ACS, and historical features that may preclude the use of fibrinolytics.

For a patient with STEMI, the goals of reperfusion are:

- First medical contact to balloon inflation within 90 minutes
- Door to drug (fibrinolytics) within 30 minutes of arrival.

The 12-lead ECG is at the center of the decision pathway in managing ischemic chest discomfort and is the only way to identify STEMI.

You can see an example ECG of an anterior STEMI below.



PREVIOUS

1 2 3

**NEXT**

CHALLENGE US




Coach



Push ► for reading content a old

Autoplay ☒ On



Self-Assessment ?


Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

# CONCURRENT ED OR CATH LAB ASSESSMENT

Below you can see an example of how to measure ST-segment elevation



**A. Inferior MI** The ST segment has no low point (it is covered or concave).

PREVIOUS

1 2 3

NEXT

CHALLENGE US ?



Coach



Push ► for reading content aloud.

Autoplay ☒ On




## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## CONCURRENT ED OR CATH LAB ASSESSMENT



B. Anterior MI

PREVIOUS

1

2

3

NEXT

CHALLENGE US









Coach



Push for reading content aloud

Autoplay On



Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## CONCURRENT ED OR CATH LAB ASSESSMENT

### Patient General Treatment

Unless allergies or contraindications exist, **consider these 4 agents** in patients with ischemic-type chest discomfort

- If  $O_2$  saturation is less than 90%, **start oxygen** at 4 L/min, titrate.
- **Aspirin** 162 to 325 mg (if not given by EMS).
- **Nitroglycerin** sublingual or translingual.
- **Morphine** IV if discomfort not relieved by nitroglycerin.
- Consider administration of P2Y<sub>12</sub> inhibitors.

Because out-of-hospital providers may have given these agents already administer initial or supplemental doses as indicated

PREVIOUS 1 2 **3**

CHALLENGE US



21% **PROGRESS:** HeartCode ACLS 2025

4h 6m left

Ahmed Othman



## Coach



Learn more here

Concurrent ED or Cath



## Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

What is the time goal for how quickly you should complete a flowchart checklist once the patient arrives in the emergency department?

You got it!



Your Answer

10 minutes



Know it

CHALLENGE US

NEXT



## Coach



Maybe this can help you?

☐ Administering Oxygen



- A patient without dyspnea has signs of acute coronary syndrome
- There are no obvious signs of heart failure
- You ~~are~~ noninvasively monitored oxygenoglobin saturation

What is the oxygen saturation threshold below which supplemental oxygen would be required?

**Not there yet...**

Your Answer  92%

Correct Answer  90%

[Learn more here](#)

☐ Administering Oxygen




Think So

CHALLENGE US

**NEXT**

## Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



Coach



Learn more here:

☐ Administering Aspirin (...



What blood component is acted upon by aspirin administration during the management of a patient with acute coronary syndromes?

You got it!



Your Answer

**Platelets**

Platelets are one of the principal and earliest participants in thrombus formation.



Know it

CHALLENGE US



Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



23% **PROGRESS** HeartCode ACLS 2025

4/15 min 1:01

Ahmed Othman

**A**



## Coach



Learn more here:

☐ Administering Aspirin (.pdf)



Which is a contraindication to the administration of aspirin for the management of a patient with acute coronary syndrome?

You got it!



Your Answer

Recent gastrointestinal bleeding



Think So

CHALLENGE US

NEXT

## Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

Coach



Push ► for reading content  
around.

Autoplay ☒ On

## ADMINISTERING ASPIRIN (ACETYLSALICYLIC ACID)

A dose of 162 to 325 mg of non-enteric-coated or chewed aspirin causes immediate and near-total inhibition of thromboxane  $A_2$  production by inhibiting platelet cyclooxygenase (COX-1).

Platelets are one of the principal and earliest participants in thrombus formation. This rapid inhibition also reduces coronary reocclusion and other recurrent events independently and after fibrinolytic therapy.



### How to Give Aspirin

If the patient has not taken aspirin and has no history of true aspirin allergy and no evidence of recent GI bleeding, give the patient aspirin (162 to 325 mg) to chew. In the initial hours of an ACS, aspirin is absorbed better when chewed than when swallowed, particularly if the patient has received morphine.

Use rectal aspirin suppositories (300 mg) for patients with nausea, vomiting, active peptic ulcer disease, or other disorders of the upper GI tract.

Aspirin is associated with a reduction in mortality for patients with ACS.

CHALLENGE US



CLOSE



Coach



Push ▶ for reading content ahead.

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## ADMINISTERING NITROGLYCERIN (GLYCERYL TRINITRATE)

Routine use of IV nitroglycerin is not indicated and has not been shown to significantly reduce mortality in STEMI. However, IV nitroglycerin is indicated and used widely in ischemic syndromes and is preferred over topical or long-acting forms because it can be adjusted in a patient with potentially unstable hemodynamics and clinical condition.

### Indications for initiating IV nitroglycerin in STEMI

- Recurrent or continuing chest discomfort unresponsive to sublingual or translingual nitroglycerin
- Pulmonary edema complicating STEMI
- Hypertension complicating STEMI

### Treatment goals using IV nitroglycerin

PREVIOUS

1

2

3

NEW

NOV

NOV

DONT GET IT

CHALLENGE US



Coach



Push ► for reading content aloud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## ADMINISTERING NITROGLYCERIN (GLYCERYL TRINITRATE)

Nitroglycerin effectively reduces ischemic chest discomfort and it has beneficial hemodynamic effects. The physiologic effects of nitrates reduce left ventricular (LV) and right ventricular (RV) preload through peripheral arterial and venous dilation.



### How to Give Nitroglycerin

Give the patient 1 sublingual nitroglycerin tablet (or translingual dose) every 3 to 5 minutes for ongoing symptoms if permitted by medical control and no contraindications exist. You may repeat the dose twice (total of 3 doses).

Administer nitroglycerin only if the patient remains hemodynamically stable: systolic blood pressure (SBP) greater than 90 mm Hg or no lower than 30 mm Hg below baseline (if known) and a heart rate of 50 to 100/min.

1

2

3

NEXT

CHALLENGE US





## Coach



Push ► for reading content aloud.

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## ADMINISTERING NITROGLYCERIN (GLYCERYL TRINITRATE)

Nitroglycerin is a venodilator. Use it cautiously or not at all in patients with inadequate ventricular preload. These situations include:

- **Inferior wall MI and RV infarction** RV infarction may complicate an inferior wall MI. Patients with acute RV infarction depend on RV filling pressures to maintain cardiac output and blood pressure. If you cannot rule out RV infarction, use caution in administering nitrates to patients with an inferior STEMI. If you confirm RV infarction by right-sided precordial leads, or if an experienced provider confirms it through clinical findings, then nitroglycerin and other vasodilators such as morphine or volume-depleting drugs (diuretics) are contraindicated as well.
- **Hypotension, bradycardia, or marked tachycardia** Avoid using nitroglycerin in patients with hypotension (SBP less than 90 mm Hg), marked bradycardia (less than 50/min), or tachycardia.
- **Recent phosphodiesterase inhibitor use** Avoid using nitroglycerin if you suspect or know that the patient has taken sildenafil or vardenafil within the previous 24 hours or tadalafil within 48 hours. These agents are generally used for erectile dysfunction or in cases of pulmonary hypertension, and in combination with nitrates, they may cause severe hypotension refractory to vasopressor agents.

PREVIOUS

1

2

3

NEXT

CHALLENGE US





Coach



Push ► for reading content aloud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## ADMINISTERING NITROGLYCERIN (GLYCERYL TRINITRATE)

### Treatment goals using IV nitroglycerin

For relief of ischemic chest discomfort:

- Titrate to effect
- Keep SBP greater than 90 mm Hg.
- Limit drop in SBP to 30 mm Hg below baseline in hypertensive patients.

For improvement in pulmonary edema and hypertension.

- Titrate to effect
- Limit drop in SBP to 10% of baseline in normotensive patients
- Limit drop in SBP to 30 mm Hg below baseline in hypertensive patients.

PREVIOUS

1

2

3

NEW

GO

DON'T GET IT

CHALLENGE US





Coach



Learn more here:

☐ Administering Nitroglycerin.



What is a physiologic effect of nitroglycerin?

You got it!



Your Answer

Reduces preload



Know it

CHALLENGE US



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



Ahmed Othman



Coach



Learn more here;

☐ Administering Nitrogly



Which clinical finding represents a contraindication to the administration of nitroglycerin?

You go



Your Answer

Confirmed right ventricular infarction



Know it

CHALLENGE US



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER





## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## ADMINISTERING MORPHINE



## Indications

Consider administering morphine for **severe chest discomfort that does not respond to sublingual or translingual nitroglycerin**, if authorized by protocol or medical control. Healthcare providers can consider giving analgesics such as morphine while monitoring the patient's blood pressure and respiratory rate. Morphine is indicated (in STEMI) when chest discomfort does not respond to nitrates.

Use morphine with caution in NSTEMI-AACS because of an association with increased mortality.

In addition, morphine may mask symptoms of myocardial ischemia and decrease absorption of important orally administered drugs (such as antiplatelets, P2Y<sub>12</sub> receptor blockers). Use morphine with caution for patients with unstable angina.

1

2

NEXT

CHALLENGE US ?



Coach



Push ► for reading content aloud.

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## ADMINISTERING MORPHINE

### ♥ Physiologic Effects

Morphine may be used to manage ACS because it

- Produces central nervous system analgesia, which reduces the adverse effects of neurohumoral activation, catecholamine release, and heightened myocardial oxygen demand
- Alleviates dyspnea
- Produces venodilation, which reduces LV preload and oxygen requirement
- Decreases systemic vascular resistance, which reduces LV afterload
- Helps redistribute blood volume in patients with acute pulmonary edema

Remember **morphine is a venodilator**. As with nitroglycerin, **use smaller doses and carefully monitor physiologic response** before administering additional doses in patients who may be **preload dependent**. If hypotension develops, administer fluids as a first line of therapy.

PREVIOUS

1

2



CHALLENGE US





## Coach



Learn more here:

☐ Administering Morphine



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

Which class of medications commonly given to patients with acute coronary syndromes may be adversely affected by morphine administration?

You got it!



Your Answer

### Oral antiplatelet medications

Morphine can decrease absorption of oral antiplatelet medications, which include clopidogrel, prasugrel, and ticagrelor.



Know it

CHALLENGE US

NEXT



## Coach



Learn more here

☐ Administering Morphine



## Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

What is a benefit of morphine when given to the management of acute coronary syndromes?

You got it!



Your Answer

Central nervous system analgesia



Know it

CHALLENGE US

NEXT

Coach



Push ▶ for reading content aloud

Autoplay ☒ On



Self-Assessment ?


Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## CLASSIFYING PATIENTS BASED ON ST-SEGMENT DEVIATION

Patients with coronary atherosclerosis may develop a spectrum of clinical syndromes that represent varying degrees of coronary artery occlusion. These syndromes include non-ST-segment elevation ACS, or NSTEMI, and ST-segment elevation myocardial infarction or STEMI. Sudden cardiac death may occur with any of these syndromes. The figure illustrates the pathophysiology of ACS.



1

NEXT

CHALLENGE US ?

Coach



Play ▶ to start the presentation

Autoplay ⏮ ⏪ ⏩ ⏭



Self-Assessment



100% 0/100



Autoplay ⏮ ⏪ ⏩ ⏭

Early plaque formation



Significant plaque formation



Plaque rupture/thrombus



Non-STEMI-ACS



STEMI



Resolution/stable angina

A Unstable plaque

B Plaque rupture

C Unstable angina

D Microemboli

E Occlusive thrombus

Coach



Push ► for reading content aloud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## CLASSIFYING PATIENTS BASED ON ST-SEGMENT DEVIATION

### STEMI

STEMI is characterized by ST-segment elevation in 2 or more contiguous leads or new left bundle branch block (LBBB)

Threshold values for ST-segment elevation consistent with STEMI are

- J-point elevation greater than 2 mm (0.2 mV) in leads  $V_2$  and  $V_3$ , 2.5 mm in men younger than 40 years, 1.5 mm in all women
- 1 mm or more in all other leads or by new or presumed new LBBB

### NSTEMI-ACS


High risk

High risk NSTEMI-ACS is characterized by ischemic ST segment depression of 0.5 mm (0.05 mV) or greater or dynamic T wave inversion with pain or

PREVIOUS

1

2



CHALLENGE US





## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## CLASSIFYING PATIENTS BASED ON ST-SEGMENT DEVIATION

## High-risk

High-risk NSTEMI-ACS is characterized by ischemic ST-segment depression of 0.5 mm (0.05 mV) or greater or dynamic T-wave inversion with pain or discomfort.

Nonpersistent or transient ST-segment elevation of 0.5 mm or greater for less than 20 minutes is also included in this category.

## Low- to intermediate-risk

Low- to intermediate-risk NSTEMI-ACS is characterized by normal or nondiagnostic changes in the ST segment or T waves that are inconclusive and require further risk stratification.

This classification includes patients with normal ECGs and those with ST-segment deviation in either direction of less than 0.5 mm (0.05 mV) or T-wave inversion of 2 mm (0.2 mV) or less.

Serial cardiac studies and functional testing are appropriate.

PREVIOUS

1

2

❌ I KNEW IT

✅ I GOT IT NOW

🔥 I'VE GOT IT

💔 I DON'T GET IT

CHALLENGE US





## Coach



Learn more here

☐ Classifying Patients Bas...



You obtain a 12-lead ECG in a patient with retrosternal chest pain.

Which electrocardiographic finding is suggestive of high-risk non-ST-segment elevation acute coronary syndrome?

You got it!



Your Answer

Dynamic T-wave inversion



Know it

CHALLENGE US



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



## Coach




Learn more here

☐ Classifying Patients Bas...



## Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

Upon reviewing a patient's 12-lead ECG, you note ST-segment elevation of 2 mm in leads II, III, and aVF.

How would you classify the electrocardiographic finding?

You got it!



Your Answer

ST-segment elevation myocardial infarction



Know it?

CHALLENGE US





Coach



Push ► for reading content  
a o l d.

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the  
right to focus on the questions



ADVANCED BEGINNER

## TREATMENT STRATEGIES FOR ACS PATIENTS



### Strategies

Patients with STEMI usually have complete occlusion of an epicardial coronary artery. Treat STEMI by providing early reperfusion therapy achieved with primary percutaneous coronary intervention (or PCI, or fibrinolytics). Reperfusion therapy opens an obstructed coronary artery with either mechanical means or drugs.

PCI performed in the cardiac catheterization laboratory after coronary angiography, allows balloon dilation and/or stent placement for an obstructed coronary artery.

Early fibrinolytic therapy or direct catheter-based reperfusion is an established standard of care for patients with STEMI who present within 12 hours after symptom onset with no contraindications.

1

2

3

4

NEXT

CHALLENGE US





Coach



Push ► for reading content aloud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## TREATMENT STRATEGIES FOR ACS PATIENTS



### Early Reperfusion Therapy

Rapidly identify patients with STEMI and use a fibrinolytic checklist to screen for indications and contraindications to fibrinolytic therapy if appropriate

The first qualified physician who encounters a patient with STEMI should interpret or confirm the 12-lead ECG, determine the risk/benefit of reperfusion therapy and direct administration of fibrinolytic therapy or activation of the PCI team. Early activation of PCI may occur with established protocols.

#### Recommended time frames

- For PCI, the goal is first medical contact to balloon inflation time of 90 minutes or less
- For patients at a non-PCI-capable hospital, time from first medical contact to device should be less than 120 minutes when considering primary PCI, but systems should strive to achieve the shortest time possible
- If fibrinolysis is the intended reperfusion, the longest acceptable ED door-to-needle time (needle time is the beginning of infusion of a fibrinolytic agent) is 30 minutes, but systems should strive to achieve the shortest time possible

PREVIOUS

1

2

3

4

NEXT

CHALLENGE US




## Coach



Push ▶ for reading content aloud.

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

# TREATMENT STRATEGIES FOR ACS PATIENTS

Rapidly identify patients with STEMI and use a fibrinolytic checklist to screen for contraindications and if contraindication is to fibrinolytic therapy — appropriate

The first qualified physician who encounters a patient with STEMI should interpret or confirm the 12-lead ECG, determine the risk/benefit of reperfusion therapy and direct administration of fibrinolytic therapy or activation of the PCI team. Early activation of PCI may occur with established protocols.

### Recommended time frames

- For PCI, the goal is first medical contact to balloon inflation time of 90 minutes or less
- For patients at a non-PCI-capable hospital, time from first medical contact to device should be less than 120 minutes when considering primary PCI but systems should strive to achieve the shortest time possible
- If fibrinolysis is the intended reperfusion, the longest acceptable ED door-to-needle time (needle time is the beginning of infusion of a fibrinolytic agent) is 30 minutes but systems should strive to achieve the shortest time possible
- Consider patients who are ineligible for fibrinolytic therapy for transfer to a PCI facility regardless of delay but prepare for a door-to-departure time of 30 minutes

PREVIOUS

1

2

3

4

NEXT

CHALLENGE US





## Coach



Push ► for reading content aloud.

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## TREATMENT STRATEGIES FOR ACS PATIENTS



## Choosing Primary PCI

The most common form of PCI is coronary angioplasty with stent placement and primary PCI is preferred over fibrinolytic administration. Many studies have shown PCI to be superior to fibrinolysis in the combined end points of death, stroke, and reinfarction for patients presenting between 3 and 12 hours after onset.

Interventional strategies for the management of STEMI are as follows.

**Primary PCI** The patient is taken to the catheterization laboratory for PCI immediately after hospital presentation.

**Rescue PCI** The patient is initially treated with fibrinolytic therapy. The patient does not show signs of reperfusion (lack of ST resolution more than 50% after 1 hour of fibrinolytic therapy administration) and therefore is referred for rescue PCI.

**Pharmacoinvasive strategy** The patient is initially treated with fibrinolytic therapy with the intention to perform coronary angiography and PCI, if appropriate.

## Considerations for the use of primary PCI

- PCI is the treatment of choice for the management of STEMI when it can be performed effectively with a first medical contact-to-balloon inflation time of 90 minutes or less by a skilled provider at a skilled PCI

PREVIOUS

1

2

3

4

NEXT

CHALLENGE US





## Coach



Push ▶ for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## TREATMENT STRATEGIES FOR ACS PATIENTS

1 hour of fibrinolytic therapy administration) and therefore is referred for rescue PCI.

Pharmacoinvasive strategy. The patient is initially treated with fibrinolytic therapy with the intention to perform coronary angiography and PCI if appropriate.

**Considerations for the use of primary PCI**

- PCI is the treatment of choice for the management of STEMI when it can be performed effectively with a first medical contact to balloon inflation time of 90 minutes or less by a skilled provider at a skilled PCI facility
- Primary PCI may also be offered to patients presenting to non-PCI-capable centers if PCI can be initiated promptly within 120 minutes after first medical contact
- For patients admitted to a non-PCI center, transferring for PCI vs administering on-site fibrinolytics may have some benefit in terms of reinfarction, stroke, and a trend to lower mortality when PCI is performed within 120 minutes after first medical contact
- PCI is also preferred in patients with contraindications to fibrinolytics and is indicated in patients with high-risk features, heart failure complicating MI, or cardiogenic shock.

PREVIOUS

1

2


3

4

NEXT

CHALLENGE US ?

Coach




Push ▶ for reading content aloud

Autoplay ☒ On

▶

Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

# TREATMENT STRATEGIES FOR ACS PATIENTS

## Using Fibrinolytic Therapy

Administer a fibrinolytic agent or "clot buster" to patients with ST-segment elevation greater than 2 mm (0.2 mV) in leads V<sub>2</sub> and V<sub>3</sub> and 1 mm or more in all other leads or by new or presumed new LBBB (leads II, aVF, leads V<sub>3</sub>, V<sub>4</sub>, leads I and aVL) without contraindications. Fibrin-specific agents achieve normal flow in about 50% of patients given these drugs.

Examples of fibrin-specific drugs are alteplase, reteplase, and tenecteplase. Streptokinase was the first fibrinolytic used widely but it is not fibrin specific.

**Considerations for the use of fibrinolytic therapy**

- in the absence of contraindications and in the presence of a favorable risk-benefit ratio, fibrinolytic therapy is one option for reperfusion in patients with STEMI and onset of symptoms within 12 hours after presentation with qualifying ECG findings and if PCCs not available within 90 minutes after first medical contact.
- in the absence of contraindications, it is also reasonable to give fibrinolytics to patients with onset of












## Coach



Push ► for reading content aloud.

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## TREATMENT STRATEGIES FOR ACS PATIENTS

- In the absence of contraindications, it is also reasonable to give fibrinolytics to patients with onset of symptoms within the prior 12 hours and ECG findings consistent with true posterior MI. Experienced providers will recognize this as a condition where ST-segment depression in the early precordial leads ( $V_1$ - $V_3$ ) is equivalent to ST-segment elevation in others. When these changes are associated with other ECG findings, it suggests a "STEMI" on the posterior wall of the heart.
- Fibrinolytics are generally not recommended for patients presenting more than 12 hours after onset of symptoms. But they may be considered if ischemic chest discomfort continues with persistent ST-segment elevation.

⊗ Do not give fibrinolytics to the following patients:

- Those who present more than 24 hours after the onset of symptoms
- Those with ST-segment depression unless a true posterior MI is suspected

PREVIOUS

1

2

3

4

NEW

NEW

NEW

DONT GET IT

CHALLENGE US





## Coach



Learn more here:

Treatment Strategies fo



What is the goal for first medical contact in balloon inflation time for patients receiving percutaneous coronary intervention?

You got it!



Your Answer

| 90 minutes




Know it

CHALLENGE US



## Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



## Coach



Learn more here:

 Treatment Strategies for



## Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

What is the longest acceptable emergency department door-to-needle time when fibrinolysis is the intended reperfusion strategy?

You got it!



Your Answer

30 minutes



Know it

CHALLENGE US





Coach



Learn more here

Treatment Strategies for...



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

What is the recommended time window after symptom onset for early fibrinolytic therapy or direct thrombolysis-based reperfusion for patients with ST-segment elevation myocardial infarction and no contraindications?

You got it!



Your Answer

Within 12 hours



Know it

CHALLENGE US


NEXT

## Coach



Push ► for reading content aloud


Autoplay ☒ On



### Introduction

A 49-year-old man says that he has had chest discomfort and excessive sweating for the past 25 minutes. The pain is not relieved with rest.

CHALLENGE US



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions




ADVANCED BEGINNER

## Coach



Push ► for reading content aloud

Autoplay ☒ Or ☐



CHOOSE THE CORRECT ANSWER

Intubate the patient immediately

Do not start oxygen

Administer albuterol nebulizer

Start oxygen at 4 L/min via nasal cannula

I KNOW IT

MINUS KNOW IT

NEED HELP

NO IDEA

## Self-Assessment ?

Adjust your difficulty estimate to the right to help us tailor your course



0% DIFFICULTY ESTIMATE

ADVANCED BEGINNER

## Coach



Push ► for reading content  
around.

Autoplay ☒ On

## Self Assessment ?

Adjust your competence estimate to the  
right to focus on the questions



ADVANCED BEGINNER

What additional questions help you determine next steps?



SELECT ALL THAT APPLY

- ☒ When was the last time you went to the doctor?
- ☒ When did the symptoms start?
- ☒ Have you had recent falls?
- ☒ Do you take any medication?
- ☒ Do you have any allergies?

I KNOW IT

FINISH KNOWING

NOT SURE

NO IDEA




## Coach



Push ► for reading content aloud.

Autoplay ☒ Or




## Self Assessment ?

Adjust your competence estimate to the right to focus on the questions



What is your interpretation of the patient's ECG tracing:



I KNOW IT

PROBABLY

NOT SURE

NO IDEA






## Coach



Push ► for reading content  
around

Autoplay ☒ On



CHOOSE THE CORRECT ANSWER

☒ Anterior ST-segment elevation myocardial infarction (STEMI)

☐ Ventricular tachycardia

☐ Posterior ST-segment elevation myocardial infarction (STEMI)

☐ Normal sinus rhythm with premature ventricular contractions

☒ I KNOW IT

☐ I THINK I KNOW

☐ I AM NOT SURE

☐ NO IDEA

## Self Assessment ?

Adjust your competence estimate to the  
right to focus on the questions



ADVANCED BCG NNER

## Coach



Push ▶ for reading content about

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the question



ADVANCED BEGINNER

With the diagnosis of STEMI, what is the most probable treatment?

CHOOSE THE CORRECT ANSWER

Admission to an intensive care unit

Release to home

Admission for observation

☒ Admission for PCI or fibrinolysis

I KNOW IT

AM NOT SURE

NOT SURE

NO IDEA

## Coach



Push ▶ for reading content aloud.

Autoplay ☒ On

## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

After you give your report to the hospital staff, advise you to proceed to the cath lab for PCI.

What is the goal for PCI when treating this patient?

CHOOSE THE CORRECT ANSWER

Door-to-needle time of 90 minutes

First medical contact-to-needle time of 30 minutes

Door-to-balloon inflation time of 30 minutes

☒ First medical contact-to-balloon inflation time of 90 minutes

I KNOW IT

NOT KNOW

CONFIDENT

NO IDEA

## Coach



Push ► for reading content  
around

Autoplay ☒ On

## Self-Assessment ?

Adjust your competence estimate to the  
right to focus on the questions



ADVANCED BEGINNER

Within the first 10 minutes, on the basis of the patient showing symptoms suggestive of myocardial ischemia, what will your first actions include?

SELECT ALL THAT APPLY

- ☐ Administer a blood thinner
- ☒ Provide prehospital notification to the receiving hospital
- ☒ Administer aspirin
- ☒ If considering prehospital fibrinolysis, use the fibrinolytic checklist
- ☒ Consider oxygen, nitroglycerin, and morphine if needed
- ☐ Administer epinephrine 1 mg IV
- ☒ Assess airway, breathing, and circulation (ABCs)
- ☐ Obtain a 12-lead electrocardiogram (ECG)

I KNOW IT

MINIKNOW

NEED HELP

NO IDEA




## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?


Adjust your competence estimate to the right to focus on the questions





ADVANCED BEGINNER


Within the first 10 minutes, on the basis of the patient showing symptoms suggestive of myocardial ischemia, what will your first actions include?


You got it

- 
 Your Answer


Provide prehospital notification to the receiving hospital
- 
 Your Answer


Administer aspirin
- 
 Your Answer

If considering prehospital fibrinolysis, use the fibrinolytic checklist
- 
 Your Answer

Assess airway, breathing, and circulation (ABCs)
- 
 Your Answer

Algorithm

[Learn more](#)
- 
 Your Answer


Obtain a 12-lead electrocardiogram (ECG)
- 
 Your Answer

Consider oxygen, nitroglycerin, and morphine if needed



I Know It

CHALLENGE US


 NEXT

## Coach



Push ► for reading content  
around

Autoplay ☒ On



His initial vital signs are HR 120/min, BP 135/88 mm Hg, RR 22/min, SpO<sub>2</sub> 87% and temperature 37.3°C.

When considering oxygen saturation, what is your next intervention?

You got it!

✓ Your Answer

Start oxygen at 4L/min via nasal cannula

Algorithm

Learn more



Knowledge

CHALLENGE US

NEXT

## Self-Assessment ?

Adjust your competence estimate to the right to focus on it, or just a hint.



ADVANCED BEG NNER

## Coach



Push ► for reading content around

Autoplay ☒ On

## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

What additional questions help you determine next steps?



SELECT ALL THAT APPLY

- ☒ Have you had recent falls?
- ☒ When was the last time you went to the doctor?
- ☒ Do you have any allergies?
- ☒ When did the symptoms start?
- ☒ Do you take any medication?

I KNOW IT

THINK I KNOW IT

NOT SURE


NO IDEA

## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

✓ Your Answer

Do you have any allergies?

✓ Your Answer

When did the symptoms start?

[Learn more](#)

✓ Your Answer

Do you take any medication?

✗ Your Answer

Have you had recent falls?

Identifying when the onset of symptoms occurred is critical in identifying and communicating to the hospital. The receiving hospital will need to determine the need and the strategy if brachytherapy or percutaneous coronary intervention or reperfusion therapy. Some prehospital providers have standing orders for reperfusion treatment.

✗ Your Answer

When was the last time you went to the doctor?

Identifying when the onset of symptoms occurred is critical in identifying and communicating to the hospital. The receiving hospital will need to determine the need and the strategy if brachytherapy or percutaneous coronary intervention or reperfusion therapy. Some prehospital providers have standing orders for reperfusion treatment.



I Know It

CHALLENGE US

NEXT





## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

What additional questions help you determine next steps?



You got it!

✓ Your Answer

Do you have any allergies?

■ Your Answer

When did the symptoms start?

[Learn more](#)

■ Your Answer

Do you take any medication?



Think So

CHALLENGE US


NEXT

## Coach




Push ► for reading content about

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions.



ADVANCED BEGINNER

Your patient continues to say that he has chest discomfort.

What is the next step? Select all that are recommended by your algorithm.

You got it!

✓ Your Answer

Nitroglycerin sublingual or translingual every 3 to 5 minutes

**Algorithm**

[Learn more](#)



Think So

CHALLENGE US


NEXT

## Coach



Push ► for reading content aloud

Autoplay ☒ On




## Self Assessment ?

Adjust your competence estimate to be right to focus on the questions



ADVANCED BEGINNER



You got it!

✓ You Answer

Anterior ST-segment elevation myocardial infarction (STEMI)

[Learn more](#)



[ Show ]

CHALLENGE US


MDG

## Coach



Push ▶ for reading content around

Autoplay ☒ On



With the diagnosis of STEMI, what is the most probable treatment?

You got it!

✓ You Answer

Admission for PCI or fibrinolysis

**Algorithm**

[Learn more](#)



1/30 play 11

CHALLENGE US

**NEXT**

## Self-Assessment ?

Adjust your competence estimate to the right to focus on the question



ADVANCED BEGINNER

## Coach



Push ► for reading content about


Autoplay ☒ On

## Self Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



✓ Your Answer

First medical contact-to-balloon inflation time of 90 minutes

**Algorithm**

[Learn more](#)



I know it

CHALLENGE US





28%

PROGRESS

To exit full screen, move mouse to top of screen or press F11

4h 17m left



Ahmed Othman



Coach



Push ► for reading content aloud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## STROKE INTRODUCTION



### Stroke Chain of Survival

The goal of stroke care is to minimize brain injury and maximize the patient's recovery. The **Stroke Chain of Survival**, described by the AHA and the American Stroke Association, or ASA, links the actions that patients, family members, and healthcare providers should take to maximize stroke recovery.

- Rapid recognition of and reaction to stroke warning signs and symptoms
- Rapid use of 9-1-1 and EMS dispatch
- Rapid EMS recognition of stroke, triage, transport, and prehospital notification to the receiving hospital
- Rapid diagnosis and treatment in the hospital



1


2

3

NEXT

CHALLENGE US







## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

# STROKE INTRODUCTION



## The 8 D's of Stroke Care

The 8 D's of Stroke Care highlight the major steps in diagnosis and treatment of stroke and key points at which delays can occur

- **Detection:** rapid recognition of stroke signs and symptoms
- **Dispatch:** early activation and dispatch of EMS by phoning 9-1-1
- **Delivery:** rapid EMS stroke identification, management, triage, transport, and prehospital notification
- **Door:** emergent ED/imaging suite triage and immediate assessment by the stroke team
- **Data:** rapid clinical evaluation, laboratory testing, and brain imaging
- **Decision:** establishing stroke diagnosis and determining optimal therapy selection
- **Drug/Device:** administration of fibrinolytic and/or endovascular therapy if eligible
- **Disposition:** rapid admission to the stroke unit or critical care unit, or emergent interfacility transfer for endovascular therapy (or EVT)

PREVIOUS

1

2

3

NEXT

CHALLENGE US







Coach



Push ▶ for reading content aloud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions




ADVANCED BEGINNER

# STROKE INTRODUCTION

## Adult Suspected Stroke Algorithm

For more information on these critical elements, see the **Adult Suspected Stroke Algorithm** below

Adult Suspected Stroke Algorithm



PREVIOUS

1

2

3

I KNOW

GOT IT NOW

THINK I GOT IT

I DIDN'T GET IT

CHALLENGE US






## Coach



Push ► for reading content  
aoud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the  
right to focus on the questions



ADVANCED BEGINNER

## MAJOR TYPES OF STROKE

**Stroke** is a general term. It refers to an **acute neurologic impairment** that follows **interruption in blood supply** to a specific area of the brain. The **major types of stroke** are **hemorrhagic** and **ischemic**.

### Hemorrhagic Stroke

**Hemorrhagic stroke** accounts for **13% of all strokes** and occurs when a **blood vessel in the brain suddenly ruptures** into the surrounding tissue.

### Ischemic Stroke

**Ischemic stroke** accounts for **87% of all strokes** and is usually caused by an **occlusion of an artery** to a region of the brain.



CHALLENGE US ?



Coach



Push ▶ for reading content aoud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions




ADVANCED BEGINNER

MAJOR TYPES OF STROKE



Area of infarction surrounding the immediate site and distal portion of brain tissue after an occlusion.



Area of ischemic penumbra (ischemic but not yet infarcted (dead) brain tissue) surrounding areas of infarction

KNOW

KNOW

KNOW

DONT GET IT

CHALLENGE US ?



28% PROGRESS HeartCode ACLS 2025

4h 19m left

TT Ahmed Othman



## Coach



What is the most common type of stroke?

You got it

Learn more here!

☐ Major Types of Stroke



Your Answer

Ischemic stroke

Ischemic strokes account for 87% of all strokes in the United States




Know it

CHALLENGE US

NEXT

## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



28% PROGRESS HeartCode ACLS 2025

4h 15m left

Ahmed Othman



## Coach



Learn more here;

☐ Major Types of Stroke



What type of stroke occurs when a blood vessel in the brain suddenly ruptures into the surrounding tissue?

You got it!



Your Answer

Hemorrhagic stroke



Know it

CHALLENGE US



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



Coach



Push ► for reading content aloud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

# WARNING SIGNS AND SYMPTOMS OF ISCHEMIC STROKE

## Signs and Symptoms

The signs and symptoms of a stroke may be **subtle**. They include

- Sudden weakness or numbness of the face, arm or leg especially on one side of the body
- Trouble speaking or understanding
- Sudden trouble seeing in one or both eyes
- Sudden trouble walking
- Dizziness or loss of balance or coordination
- Sudden severe headache with no known cause
- Sudden confusion

CHALLENGE US ?



Coach



Learn more here

☐ Warning Signs and Sym...



Which is a sign of stroke?

You got it!



Your Answer

Trouble speaking



Know it

CHALLENGE US

NEXT

Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



31% PROGRESS: HeartCode ACLS 2025

4h 15m left

Ahmed Othman

A



Coach



Learn more here

Warning Signs and Sym.



Which is a symptom of stroke?

You got it!



Your Answer

Sudden trouble seeing



Know it

CHALLENGE US

NEXT

Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER






## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

# STROKE ASSESSMENT TOOLS FOR EMS

The AHA recommends that all EMS personnel be trained to recognize stroke by using a validated, abbreviated out-of-hospital neurologic evaluation tool such as the Cincinnati Prehospital Stroke Scale (CPSS) or the Los Angeles Prehospital Stroke Screen.

The CPSS identifies stroke on the basis of 3 physical findings

- **Facial droop** (have the patient smile or try to show teeth)
- **Arm drift** (have the patient close eyes and hold both arms out, with palms up;
- **Abnormal speech** (have the patient say, "You can't teach an old dog new tricks")

By using the CPSS, medical personnel can evaluate the patient in less than 1 minute. The presence of 1 finding on the CPSS has an estimated probability of 72% when scored by prehospital providers.

You can see a table of normal and abnormal findings in the CPSS by clicking on the CPSS button below

CPSS

LINK NEW

OPEN NOW

LINK NEW

DON'T GET IT

CHALLENGE US



## Test

## Findings

**Facial droop:** have the patient show teeth or smile

**Normal**—both sides of the face move equally

**Abnormal**—one side of the face does not move as well as the other side

**Arm drift:** patient closes eyes and extends both arms straight out, with palms up, for 10 seconds

**Normal**—both arms move the same or both arms do not move at all (other findings, such as pronator drift, may be helpful)

**Abnormal**—one arm does not move or one arm drifts down compared with the other

**Abnormal speech:** have the patient say, "you can't teach an old dog new tricks"

**Normal**—patient uses correct words with no slurring

**Abnormal**—patient slurs words, uses the wrong words, or is unable to speak

**Interpretation:** if any 1 of these 3 signs is abnormal the probability of a stroke is 72%

Modified from Kothari RU, Pancioli A, Liu T, Broderick J. Cincinnati Prehospital Stroke Scale: reproducibility and validity. *Ann Emerg Med.* 1999;33(4):373-378.

With permission from Elsevier.

CLOSE



31% **PROGRESS:** HeartCode ACLS 2025

4h 16m left

Ahmed Othman



## Coach



Learn more here:

Stroke Assessment Tool



## Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

What validated, abbreviated out-of-hospital neurologic evaluation tool contains 3 components: the facial droop, arm drift, and abnormal speech tests?

You got it!



Your Answer

Cincinnati Prehospital Stroke Scale



Know it

CHALLENGE US

NEXT



Coach



Learn more here:

[Stroke Assessment Tool.](#)



What is the estimated probability of the Cincinnati Prehospital Stroke Scale with 1 abnormal finding when scored by prehospital providers?

You got it!

Your Answer: 72%



Know it

CHALLENGE US

Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



## Coach



Push ► for reading content aloud

Autoplay ☒ On

## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

# CRITICAL EMS ASSESSMENTS AND ACTIONS

To help identify which hospital is most appropriate for the stroke patient, EMS should also perform a severity screen to determine if there is a large-vessel occlusion (LVO). As always, EMS should follow local stroke protocols.



### After Recognizing the Stroke

- Assess ABCs; give oxygen if needed
- Initiate stroke protocol
- Perform physical exam
- Perform validated prehospital stroke screen and severity tool
- Establish time of symptom onset (last known normal)
- Triage to most appropriate stroke center
- Check glucose; treat if indicated
- Provide prehospital notification on arrival; transport to brain imaging suite

1

2

3

**NEXT**

CHALLENGE US






Coach



Push ► for reading content aloud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## CRITICAL EMS ASSESSMENTS AND ACTIONS

### After Recognizing the Stroke

- Assess ABCs, give oxygen if needed
- Initiate stroke protocol
- Perform physical exam
- Perform validated prehospital stroke screen and severity tool
- Establish time of symptom onset (last known normal)
- Triage to most appropriate stroke center
- Check glucose; treat if indicated
- Provide prehospital notification on arrival, transport to brain imaging suite

Both out-of-hospital and in-hospital medical personnel should provide supplemental oxygen to hypoxemic stroke patients (those whose oxygen saturation is 94% or less) or patients for whom oxygen saturation is unknown.

1

2

3

END

CHALLENGE US






Coach



Push ► for reading content aloud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## CRITICAL EMS ASSESSMENTS AND ACTIONS

### Examples of stroke assessment tools

- Cincinnati Prehospital Stroke Scale (CPSS, FAST)
- Los Angeles Prehospital Stroke Screen (LAPSS)
- Melbourne Ambulance Stroke Screen (MASS)
- Miami Emergency Neurologic Deficit Score (MENDS)
- Recognition of Stroke in the Emergency Room Score (ROSIER)

### Stroke severity scores

- National Institutes of Health Stroke Scale (NIHSS)
- Shortened National Institutes of Health Stroke Scale 5 and 8 (sNIHSS 5 and sNIHSS 8)
- Cincinnati Prehospital Stroke Severity Screen (CPSSS)
- Field Assessment Stroke Triage for Emergency Destination (FAST-ED)
- Los Angeles Motor Scale (LAMS)

PREVIOUS

1

2

3

NEXT

CHALLENGE US





Coach



Push ▶ for reading content aloud

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## CRITICAL EMS ASSESSMENTS AND ACTIONS

- Melbourne Ambulance Stroke Screen (MASS)
- Miami Emergency Neurologic Deficit Score (MENDS)
- Recognition of Stroke in the Emergency Room Score (ROSIER)

### Stroke severity scores

- National Institutes of Health Stroke Scale (NIHSS)
- Shortened National Institutes of Health Stroke Scale 5 and 8 (sNIHSS-5 and sNIHSS-8)
- Cincinnati Prehospital Stroke Severity Screen (CPSSS)
- Field Assessment Stroke Triage for Emergency Destination (FAST-ED)
- Los Angeles Motor Scale (LAMS)
- Rapid Arterial Occlusion Evaluation score (RACE)
- 3-Item Stroke Scale (3ISS)

PREVIOUS

1 2 3

NEXT

CHALLENGE US








Coach



Push ► for reading content aloud

Autoplay ☒ On



Self-Assessment ?


Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

# CRITICAL EMS ASSESSMENTS AND ACTIONS

Emergency Medical Services Acute Stroke Routing



PREVIOUS 1 2 **3**



CHALLENGE US ?



Coach

Learn more here:

☐ Critical EMS Assessment



Which stroke severity tool that helps EMS differentiate large-vessel occlusion stroke from non-large-vessel occlusion stroke?

You got it!

Your Answer

Los Angeles Motor Scale

Know it

CHALLENGE US





33% **PROGRESS** HeartCode ACLS 2025

4h 12m left



Ahmed Othman



## Coach



Push ► for reading content  
aoud

Autoplay ☒ On

What is the primary advantage of using a stroke severity tool?

You got it!



Your  
Answer

It helps identify large-vessel occlusion stroke.



Know it

CHALLENGE US






## Coach



Push ► for reading content aloud

Autoplay ☒ On



## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

# STROKE CENTERS AND STROKE UNITS

Evidence indicates a benefit from triage of stroke patients directly to designated certified stroke centers. Currently, **4 levels of stroke certification exist**, and certification is given on the basis of a hospital's specific capabilities.



### Acute Stroke Ready Hospital

Acute Stroke Ready Hospitals typically serve **rural and under-resourced areas**. Emergent identification and treatment of patients with alteplase, when indicated, is typically facilitated by telemedicine to provide access to acute neurologic expertise. Typically, **patients are later transferred for admission to a stroke unit or for a higher level of care**, as indicated.



### Primary Stroke Center

The Primary Stroke Center is the **cornerstone of stroke systems of care**. These centers comprise a wide range of hospitals **able to quickly identify stroke patients**, provide alteplase therapy if indicated, and **admit patients to a dedicated stroke unit**. Roughly half of all stroke patients in the United States receive care in a Primary Stroke Center.

■ I KNOW

■ I THINK I KNOW

■ I DON'T KNOW

■ I DON'T GET IT

CHALLENGE US





33% PROGRESS HeartCode ACLS 2025

4h 13m left

TT Ahmed Othman



Coach



Push ► for reading content aloud.

Autoplay ☒ On



Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## STROKE CENTERS AND STROKE UNITS

The Primary Stroke Center is the cornerstone of stroke systems of care. These centers comprise a wide range of hospitals able to quickly identify stroke patients, provide alteplase therapy if indicated, and admit patients to a dedicated stroke unit. Roughly half of all stroke patients in the United States receive care in a Primary Stroke Center.



### Thrombectomy-Capable Stroke Center

The Thrombectomy-Capable Stroke Center certification was jointly created by the AHA and the Joint Commission to recognize stroke centers that meet the same high-quality standards as a primary stroke center but are also capable of providing endovascular therapies for patients with LVO. The Thrombectomy-Capable Stroke Center designation was created to recognize these EVT-capable facilities in areas where a Comprehensive Stroke Center was not available.



### Comprehensive Stroke Center

Hospitals achieving Comprehensive Stroke Center certification are capable of managing all forms and severities of stroke, both ischemic and hemorrhagic, and can provide 24/7 access to specialty care, such as neurosurgery, endovascular therapy, and neurocritical care. A Comprehensive Stroke Center typically serves as the hub of a regional stroke system of care, providing receiving capabilities for transferred patients and providing feedback and education for transferring sites.



CHALLENGE US ?



34%

PROGRESS: HeartCode ACLS 2025

—4h 12m left

TT Ahmed Othman

A



## Coach



Learn more here:

Stroke Centers and Stro...



What is the most appropriate destination for patients with suspected acute ischemic stroke?

You got it!



Your  
Answer

Certified stroke center




I Know It

CHALLENGE US

NEXT

## Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



34% **PROGRESS:** HeartCode ACLS 2025

~4h 12m left



Ahmed Othman



## Coach



Learn more here:

Stroke Centers and Stro...



What is the highest level of stroke center certification?

You got it!



Your  
Answer

| Comprehensive Stroke Center



I Know It.

CHALLENGE US

NEXT

## Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER



34%

PROGRESS: HeartCode ACLS 2025

-4h 53m left

Ahmed Othman

A



Coach



Push ► for reading content aloud.

Autoplay ☒ On

Self-Assessment ?

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

## EMS TRANSPORT AND ALERTING THE RECEIVING FACILITY



### Advantages of EMS Transport

- Emergency medical dispatchers play a critical role in timely treatment of potential stroke by
  - **Identifying** possible stroke patients
  - Providing **high-priority** dispatch
  - **Instructing** bystanders in lifesaving CPR skills or other supportive care if needed while EMS providers are on the way
- Responding providers can **assess ABCs** and **give oxygen** as needed.
- EMS personnel can **initiate stroke protocol**, **perform a physical exam**, **establish time of symptom onset (last known normal)**, and **check glucose** and **treat** if indicated.
- EMS can **triage to the most appropriate stroke center** on the basis of a validated prehospital stroke screen, stroke severity tool, and on patient characteristics following regional destination protocols.
- EMS can **provide prehospital notification**, enabling the hospital to prepare to evaluate and manage the patient more efficiently, and on arrival transport to the brain imaging suite.

I KNEW

GOT IT NOW

THINK I GOT IT

I DON'T GET IT

CHALLENGE US







Coach



Learn more here:

☐ EMS Transport and Aler...



Self-Assessment

Adjust your competence estimate to the right to focus on the questions



ADVANCED BEGINNER

Which is an advantage of EMS transport to a stroke hospital for a patient with suspected acute ischemic stroke?

You got it!



Your Answer

Responding providers can stabilize critical issues.



I Know It

CHALLENGE US

NEXT